

*This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.*



**ISSN: 2167-0390**

## **Vitamins & Trace Elements**

**The International Open Access  
Vitamins & Trace Elements**

### **Executive Editors**

**Mary Harris**

Colorado State University, USA

**Karen S. Kubena**

Texas A&M University, USA

**Duo Li**

Zhejiang University, China

**Des R. Richardson**

University of Sydney, Australia

**Raymond R. Hyatt**

Tufts University School of Medicine, USA

**Available online at:** OMICS Publishing Group ([www.omicsonline.org](http://www.omicsonline.org))

This article was originally published in a journal by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/2167-0390.1000108>

# Antioxidant Substances and Trace Element Content in Macroalgae from a Subtropical Lagoon in the West Coast of the Baja California Peninsula

Paola A. Tenorio Rodríguez<sup>1</sup>, LC Méndez-Rodríguez<sup>1</sup>, E Serviere-Zaragoza<sup>2</sup>, T O'Hara<sup>3</sup> and T Zenteno-Savín<sup>1\*</sup>

<sup>1</sup>Planeación Ambiental y Conservación, Centro de Investigaciones Biológicas del Noroeste, S.C., La Paz, Baja California Sur, México

<sup>2</sup>Ecología Pesquera, Centro de Investigaciones Biológicas del Noroeste, S.C., La Paz, Baja California Sur, México

<sup>3</sup>Department of Biology and Wildlife, University of Alaska Fairbanks, Fairbanks, USA

## Abstract

Macroalgae play an important role in the ecology of the marine environment. They form the base of the food webs, and represent the major food source for a variety of organisms. In order to assess their potential nutritional value, the antioxidant and trace element content (vitamin C, total polyphenols, zinc, iron, copper, selenium, cadmium and lead) of eight macroalgae species, three red (*Hypnea spinella*, *Gracilaria textorii* and *G. vermiculophylla*), four green (*Caulerpa sertularioides*, *Codium simulans*, *C. amplivesiculatum* and *Ulva lactuca*) and one brown (*Dictyota flabellata*) macroalgae, were determined. The concentration ranges found were as follows: zinc, 19.1-7.4; iron, 638.4-89.2; copper, 3.9-0.9; selenium, 0.32-10; cadmium, 4.33-0.02, and lead 8.8-0.39 mg kg<sup>-1</sup> dry weight. Green macroalgae showed higher concentrations of iron and zinc. Total polyphenol content ranged from 29.6 to 70.3 mg 100 g<sup>-1</sup> dry weight; brown macroalgae showed higher polyphenol concentration. *C. simulans* and *C. amplivesiculatum* had higher vitamin C content (3.16 ± 0.52 mg g<sup>-1</sup> and 6.08 ± 0.69 mg g<sup>-1</sup> dry weight, respectively) than the other species. Comparison with several commonly consumed vegetables and fruits suggest that these macroalgae are likely a good alternative source of antioxidants and micronutrients for consumers.

**Keywords:** Antioxidants; Macroalgae; Micronutrients; Polyphenols; Trace elements; Vitamin C

**Abbreviations:** ROS: Reactive oxygen species; LDL: Low density lipoproteins; O<sub>2</sub><sup>•-</sup>: Superoxide radical; H<sub>2</sub>O<sub>2</sub>: Hydrogen peroxide; OH<sup>•</sup>: Hydroxyl radical; <sup>1</sup>O<sub>2</sub>: Singlet oxygen; Cu: Copper; Fe: Iron; Pb: Lead; Cd: Cadmium; Zn: Zinc; Se: Selenium; GAE: Gallic acid equivalents; FAO/WHO: Food and Agriculture Organization/World Health Organization; RNI: Recommended nutrient intake

## Introduction

In many countries, fresh marine macroalgae are used as food by coastal communities, and considered as a traditional food item due to their nutritional value and characteristic taste [1-3]. Macroalgae are valuable sources of proteins, polysaccharides, and fiber; but they are also rich in antioxidants and micronutrients, such as vitamins and trace elements [4-6].

Antioxidant vitamins and trace elements are usually obtained from the diet, since some organisms are unable to synthesize them. The beneficial effects of antioxidants are due to their capacity to scavenge and neutralize reactive oxygen species (ROS) [7]. An excessive ROS production and/or low antioxidant defense can cause oxidative damage to biomolecules, such as proteins, lipids and DNA [8,9]. Antioxidants may reduce ROS production by scavenging free radicals through various mechanisms [7,10]. Some trace elements contribute to the function of endogenous antioxidant enzymes by acting as cofactors [11,12]. Most polyphenols can act as chain breakers or radical scavengers and prevent the oxidation of low density lipoproteins (LDL) [12,13]. Vitamin C is a scavenger against superoxide radical (O<sub>2</sub><sup>•-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), hydroxyl radical (OH<sup>•</sup>), and singlet oxygen (<sup>1</sup>O<sub>2</sub>), which would otherwise react to form lipid peroxides; vitamin C also reduces the tocopheroxyl radical formed by interaction of  $\alpha$ -tocopherol with lipid peroxides in cell membranes [7,14]. This demonstrates the wide variety of reactive intermediates vitamin C interacts with, to maintain the function of cellular components.

Marine macroalgae are a potentially good source of micronutrients and may be beneficial for human health, given their reportedly high vitamin and trace element content [1,15,16]. Most published studies

on macroalgae are focused on ecological aspects and chemical composition (protein, carbohydrate, ash and calories). However, little is known about macroalgae antioxidant vitamins and micronutrient composition [5,17,18]. The objective of this study was to assess the antioxidant vitamin and trace element content of eight macroalgae species, from a subtropical and unpolluted lagoon in the west coast of the Baja California peninsula. Lead and cadmium concentrations were also assessed, given their toxicological potential. This information may be useful in the search for alternative and supplementary food items in the Baja California peninsula.

## Materials and Methods

Eight species of macroalgae belonging to the Chlorophyta, Rhodophyta and Ochrophyta Pheophyceae divisions were collected from Bahía Magdalena, Baja California Sur, Mexico (24°15' N and 25°20' N and 111°30' W and 112°15' W). Bahía Magdalena is a shallow lagoon with high productivity resulting from seasonal marine upwelling [19]. Macroalgae were collected in November of 2009, February, April and June of 2010. Samples were carried on ice, stored in black bags to the laboratory, where macroalgae were washed with filtered, cold seawater to remove all epiphytes and other debris. The samples used for trace element analyses were dried in an oven at 60°C for 48 h. The dried samples were then ground into a fine powder using a coffee grinder, and stored until analyzed. Dried samples destined for copper (Cu), iron (Fe), lead (Pb), cadmium (Cd) and zinc (Zn) analyses were digested in

**\*Corresponding author:** T Zenteno-Savín, Planeación Ambiental y Conservación, Centro de Investigaciones Biológicas del Noroeste, S.C., Instituto Politécnico Nacional 195, Playa Palo de Santa Rita Sur, La Paz, Baja California Sur, C.P. 23096, Mexico, E-mail: [tzenteno04@cibnor.mx](mailto:tzenteno04@cibnor.mx)

**Received** November 12, 2012; **Accepted** November 24, 2012; **Published** November 26, 2012

**Citation:** Tenorio-Rodríguez PA, Méndez-Rodríguez LC, Serviere-Zaragoza E, O'Hara T, Zenteno-Savín T (2013) Antioxidant Substances and Trace Element Content in Macroalgae from a Subtropical Lagoon in the West Coast of the Baja California Peninsula. *Vitam Trace Elem* 2:108. doi:10.4172/2167-0390.1000108

**Copyright:** © 2013 Zenteno-Savín T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

acid-washed Teflon tubes, with concentrated nitric acid (HNO<sub>3</sub>) in a microwave oven (Mars 5X, CEM, Matthew, NC, USA), and analyzed by atomic absorption (GBC Scientific equipment, AVANTA, Dandegong, Australia), using an air-acetylene flame. Selenium (Se) determinations were performed at the Wildlife Toxicology Laboratory, University of Alaska Fairbanks. Samples were digested twice, first with HNO<sub>3</sub>, and then with HCl (37%) in a microwave (3000 Microwave Multiwave Sample Preparation System, Anton Para, Austria). Se concentration was analyzed by atomic absorption spectrometry (AAnalyst 800 PerkinElmer Instruments, Shelton, CT, USA), with a flow injection system, according to the procedure described by Knott et al. [20] and Barrera-García et al. [21]. The results are expressed in mg kg<sup>-1</sup> dry weight (d.w.).

The vitamin C (L-ascrobic acid) content of macroalgae was determined by high-performance liquid chromatography (HPLC, Waters, Milford, MA, USA) [22,23]. Each sample (0.1 g) was homogenized with metaphosphoric acid (3% w/v), incubated for 40 min at 4°C in the dark and centrifuged for 15 min at 23,985 g at 4°C. Samples were then filtered through a 0.45 µm filter prior to injection. Ten µL of each sample were injected into the HPLC. The mobile phase was a mixture of water, sulfuric acid (pH 2.4), and acetonitrile (100%, HPLC grade). The flow rate was 1 mL min<sup>-1</sup>, and the detection wavelength was set at 245 nm. The results are expressed in mg vitamin C g<sup>-1</sup> d.w.

The Folin-Ciocalteu colorimetric method, with modifications, was used to quantify total polyphenol concentration in macroalgae samples [24,25]. Briefly, 1 g of fresh sample was ground with a mixture of water:methanol:acetone (2:3:5 v/v), using a mortar and pestle over ice. Samples were incubated in a water bath at 65°C with agitation for 1 h. Extracts were centrifuged at 15,292 g for 10 min, and the supernatant was collected. Na<sub>2</sub>CO<sub>3</sub> was added and samples were incubated for 1 h at room temperature. The absorbance at 750 nm was recorded in a microplate reader (BioRad™ 550, Hercules, CA, USA), and compared to a gallic acid calibration curve. The results are expressed as gallic acid equivalents (GAE) in mg 100 g<sup>-1</sup> fresh weight (f.w.).

Normality and homogeneity of variance were determined by using Shapiro-Wilks and Bartlett tests. All variables were log-transformed in order to normalize the data. Differences between groups and species were analyzed by using ANOVA or Student t-tests, with Tukey post-hoc tests [26]. Results are expressed as mean ± SE, and differences were considered statistically significant at p<0.05 level. Statistica 8.0 (StatSoft Inc. Tulsa, OK, USA) software was used to perform all statistical analyses.

## Results

The trace element content of eight macroalgae species is shown in table 1. Fe concentration in all macroalgae species studied ranged from 89.2 to 638.4 mg kg<sup>-1</sup> d.w. Green macroalgae had the highest Fe content (p=0.013), with *C. simulans* and *C. amplivesiculatum* showing the highest values. Zn levels ranged from 7.4 mg kg<sup>-1</sup> d.w. in *H. spinella* to 19.1 mg kg<sup>-1</sup> d.w. in *U. lactuca*. Cu concentrations ranged from 0.9 mg kg<sup>-1</sup> in *H. spinella* to 3.9 mg kg<sup>-1</sup> in *C. sertularioides*. Se, Cd and Pb levels ranged between 0.10-0.32 mg kg<sup>-1</sup>, 0.01-4.33 mg kg<sup>-1</sup>, and 8.8-0.54 mg kg<sup>-1</sup>, respectively (Table 1). No differences in trace element content were found between groups or among species.

The mean content of total polyphenols in all macroalgae species studied, ranged from 29.6 to 70.3 mg of GAE 100 g<sup>-1</sup> w.w. (Figure 1). Total polyphenol content was higher in green and brown macroalgae, compared to red macroalgae (p=0.00002). In general, the brown macroalga *D. flabellata* and the green macroalga *C. sertularioides* had higher total polyphenol concentrations, in comparison with the rest of the species (70.3 ± 7.6 and 68.6 ± 6.03 mg of GAE 100 g<sup>-1</sup> w.w., respectively). *U. lactuca* had the lowest total polyphenol content (29.6 ± 1.5 mg of GAE 100 g<sup>-1</sup> w.w.).

Vitamin C concentration in eight macroalgae species from Bahia Magdalena, Baja California Sur is shown in figure 2. Vitamin C content in all macroalgae species analyzed ranged from 1.24 to 6.08 mg g<sup>-1</sup> d.w. Vitamin C concentration was higher in green and brown macroalgae, compared to red macroalgae (p=0.002). The highest vitamin C concentration was found in the green macroalga *C. amplivesiculatum* (6.08 ± 0.69 mg g<sup>-1</sup> d.w.) (p=0.001), while the red macroalga *H. spinella* had the lowest vitamin C content (1.24 ± 0.16 mg g<sup>-1</sup> d.w.).

## Discussion

The majority of the essential minerals and trace elements needed for human nutrition can be found in macroalgae; therefore, macroalgae could be regarded as a valuable resource. Trace elements such as Fe, Cu and Se are considered essential for biological processes, including growth, reproduction, hormone metabolism and antioxidant defense [27-29]. Cd and Pb are a potential health hazard and are commonly measured, as indicators of environmental pollution [30,31]. All macroalgae species analyzed in the present study showed lower concentrations of Cd than previously reported (0.71 mg kg<sup>-1</sup> [32]; 1.9 mg kg<sup>-1</sup> [31]; 3.70 mg kg<sup>-1</sup> [33]; 4.8 mg kg<sup>-1</sup> [34]). The mean Pb concentration in macroalgae species analyzed was 3.46 mg kg<sup>-1</sup>. Similar Pb concentrations were reported for macroalgae (3.50 mg kg<sup>-1</sup> [33], and

Species	n	Zn	Fe	Cu	Se	Cd	Pb
<b>Chlorophyta</b>							
<i>Caulerpa sertularioides</i> (S.G. Gmelin) M.A. Howe	5	16.8 ± 3.9	420.9 ± 107.9	3.9 ± 1.84	0.32 ± 0.09	0.38 ± 0.18	2.71 ± 1.01
<i>Codium amplivesiculatum</i> Setchell & N.L. Gardner	3	17.1 ± 4.9	638.4 ± 176.4	2.2 ± 0.66	0.19 ± 0.07	0.02 ± 0.00	0.39 ± 0.28
<i>Codium simulans</i> Setchell & N.L. Gardner	5	18.2 ± 6.5	447.6 ± 176.7	1.7 ± 0.52	0.19 ± 0.01	1.11 ± 0.39	8.28 ± 2.44
<i>Ulva lactuca</i> Linnaeus	3	19.1 ± 7.5	213.1 ± 72.3	2.3 ± 2.2	0.11 ± 0.04	0.02 ± 0.01	1.68 ± 0.46
<b>Ochrophyta-Phaeophyceae</b>							
<i>Dictyota flabellata</i> (F.S.Collins) Setchell & N.L.Gardner*	2	14.0	560.1	2.0		4.33	8.8
<b>Rhodophyta</b>							
<i>Gracilaria vermiculophylla</i> (Ohmi) Papenfus	7	12.2 ± 4.8	190.3 ± 51.5	1.0 ± 0.41	0.10 ± 0.02	0.10 ± 0.09	2.25 ± 1.05
<i>Gracilaria textorii</i> (Suringar) De Toni	4	15.8 ± 4.3	89.2 ± 24.4	1.2 ± 0.18	0.15 ± 0.01	1.09 ± 0.54	1.47 ± 0.77
<i>Hypnea spinella</i> (C. Agardh) Kützting	3	7.4 ± 0.21	409.6 ± 27.2	0.9 ± 0.05	0.10 ± 0.01	1.01*	2.1*


Mean values ± standard error

\*Standard error not calculated


**Table 1:** Trace element concentrations (mg kg<sup>-1</sup>) in macroalgae species from Bahia Magdalena, Baja California Sur, Mexico.

2.84 mg kg<sup>-1</sup> [35]); however, Mok et al. [36] reported lower Pb levels (0.89 mg kg<sup>-1</sup>). Se content in macroalgae has been reported to be in the range of 0.1-6 µg g<sup>-1</sup> [32], which is similar to what was found in the present study. Concentrations of trace elements similar or lower than those found in eight species of macroalgae in this study have been reported for vegetables: 323 µg Fe g<sup>-1</sup> in lettuce, 182.8 µg Fe g<sup>-1</sup> in parsley, 166.4 µg Fe g<sup>-1</sup> in spinach [37], 1.02-0.33 mg Cu kg<sup>-1</sup> in spinach, and 1.11-0.51 mg Cu kg<sup>-1</sup> in tomato [38]. The concentrations of these elements in macroalgae depend on the surrounding environment. Natural enrichment of trace elements in the Baja California peninsula has been suggested to occur due to natural phenomena, such as currents, upwelling and various geological sources [39,40].

Concentrations of vitamin C in most of the analyzed macroalgae species in this study were similar or higher than those reported for congeneric species in Hawaii (3.0, 2.2, and 1.3 mg g<sup>-1</sup> d.w. in *Ulva*



**Figure 1:** Total polyphenol content (mg 100 g<sup>-1</sup> fw) in macroalgae species from Bahia Magdalena, Baja California Sur, Mexico, grouped by division. Data are presented as mean ± SE.



**Figure 2:** Vitamin C content (mg g<sup>-1</sup> dw) in macroalgae species from Bahia Magdalena, Baja California Sur, Mexico, grouped by division. Data are presented as mean ± SE.

Micronutrient	Recommended daily intake	Reference
Fe	48 mg	Joint FAO/WHO Expert Committee on Food Additives, 1999
Cu	3 mg	Joint FAO/WHO Expert Committee on Food Additives, 1999
Zn	60 mg	Joint FAO/WHO Expert consultation 1998
Se	29 µg	Joint FAO/WHO Expert consultation 1998
Cd	70 µg	Joint FAO/WHO Expert Committee on Food Additives, 1999
Pb	214 µg	Joint FAO/WHO Expert Committee on Food Additives, 1999
Vitamin C	45 mg	Joint FAO/WHO Expert consultation 1998

**Table 2:** Recommended daily dose of trace elements and antioxidants.

*flexuosa* (as *Enteromorpha flexuosa*), *Ulva fasciata* and *Monostroma oxyspermum*, respectively, [5,41], along the coast of India (0.00244 mg g<sup>-1</sup> d.w. in *Ulva reticulata*) [42], or the central part of the Baja California Peninsula (3.4 mg g<sup>-1</sup> d.w. in *Eisenia arborea*) [43]. However, the two *Codium* species analyzed in this study had higher vitamin C content (mean of 4.62 mg g<sup>-1</sup> d.w.) than the values reported for other green macroalgae species [5,44]. Overall, vitamin C concentrations in the macroalgae species analyzed in this study were comparable with those reported in guava (1.44 mg g<sup>-1</sup>), papaya (1.08 mg g<sup>-1</sup>) and red peppers (1.51 mg g<sup>-1</sup>), and were higher than in oranges (0.67 mg g<sup>-1</sup>), or broccoli (0.89 mg g<sup>-1</sup>) [45,46]. The intake of vitamin C is essential since the human body is unable to synthesize it. It is effective in ameliorating symptoms of the common cold; moreover, evidence shows that vitamin C reduces mortality from heart disease, and can prevent oxidative stress in the process of immune response [47,48].

Total poly phenol content in the brown macroalga *D. flabellata* and in the green macroalga *C. sertularioides* was higher than in other species analyzed in this study. Previous studies similarly reported that polyphenol content in brown marine algae is higher than that in red marine algae [49,50]. Phenolic compounds are widely distributed in all macroalgae. These compounds are good antioxidants, since they can act as metal chelators and ROS scavengers trapping the lipid alkoxyl radical preventing lipid peroxidation. The beneficial effects of polyphenols on human health and their contribution to protect against chronic diseases, such as neurodegenerative disorders and cancer, have been extensively documented [51,52]. In this study, total polyphenol content in the macroalgae species analyzed was higher than that reported for several vegetables and fruits, such as avocado (3.6 mg 100 g<sup>-1</sup> f.w.), kiwifruit (28.1 mg 100 g<sup>-1</sup> f.w.), broccoli (98.9 mg 100 g<sup>-1</sup> f.w.), pear (69.2 mg 100 g<sup>-1</sup> f.w.), mango (68.1 mg 100 g<sup>-1</sup> f.w.), and tomato (13.7 mg 100 g<sup>-1</sup> f.w.) [53].

In order to evaluate the food safety of the analyzed samples, regulation in other countries was consulted, since Mexico has not yet adopted official trace element limits regarding macroalgae used for human consumption. Cd concentration in *G. textorii*, *H. spinella*, *C. simulans* and *D. flabellata* and Pb content in *C. simulans* and *D. flabellata* exceeded the permissible limit, considered by the French Legislation (0.5 mg Cd kg<sup>-1</sup> and 5 mg Pb kg<sup>-1</sup> d.w.). However, the Food and Agriculture Organization/World Health Organization (FAO/WHO) [54-56] have jointly recommended daily doses for trace elements and vitamins, based on the body weight of an average adult (60 kg body weight). Recommended nutrient intake (RNI) values are shown in table 2. The Committee on Medical Aspects of Food and Nutrition Policy [57], suggested macroalgae consumption of 8 g per day is a typical daily portion consumed in the Asian cuisine. Considering these references, and the maximum content of Pb and Cd detected in *D. flabellata* (8.8 mg kg<sup>-1</sup> and 4.33 mg kg<sup>-1</sup> d.w., respectively), these macroalgae would



contribute 33% Pb and 49% Cd of the recommended daily dose by FAO/WHO. These contributions from a single macroalgae species could be considered to be high. However, trace element levels in all the macroalgae species analyzed in this study are below the permissible limits established by FAO/WHO.

The trace element and antioxidant content found in eight species of macroalgae in Bahía Magdalena, Baja California Sur, suggest that these species are safe and potentially exploitable for human consumption, and that macroalgae could be an alternative source of antioxidants and trace elements to help meet the recommended daily intake of some trace elements and vitamin C. However, in brown macroalgae, the content of elements, such as Cd and Pb should be monitored to ensure chemical safety.

#### Acknowledgments

Authors acknowledge financial support from CIBNOR (PC2.0, PC0.10), and the assistance of H. Berviera Leon and J. Angulo Calvillo in the field, A. Mazariegos in macroalgae identification, and N. Olguín-Monroy, B. Acosta, G. Peña Armenta, O. Lugo-Lugo and J.M. Castellini in quantitative analysis.

#### References

- Chapman VJ, Chapman DJ (1980) *Seaweeds and their uses*. (3<sup>rd</sup> edition), London & New York: Chapman & Hall, USA.
- Rupérez P (2002) Mineral content of edible marine seaweeds. *Food Chem* 79: 23-26.
- MacArtain P, Gill CI, Brooks M, Campbell R, Rowland IR (2007) Nutritional value of edible seaweeds. *Nutr Rev* 65: 535-543.
- Darcy-Vrillon B (1993) Nutritional aspects of the enveloping use of marine macro algae for the human industry. *Int J Food Sci Nutr* 44: 23-35.
- McDermid KJ, Stuercke B (2003) Nutritional composition of edible Hawaiian seaweeds. *J Appl Phycol* 15: 513-524.
- Mamatha BS, Namitha KK, Senthil A, Smitha J, Ravishankar GA (2007) Studies on use of *Enteromorpha* in snack food. *Food Chem* 101: 1707-1713.
- Halliwell B, Gutteridge JMC (1985) *Free Radicals in Biology and Medicine*. Clarendon Press, New York, USA.
- Radák Z, Chung HY, Goto S (2005) Exercise and hormesis: oxidative stress-related adaptation for successful aging. *Biogerontology* 6: 71-75.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, et al. (2007) Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 1: 44-84.
- Vertuani S, Angusti A, Manfredini S (2004) The antioxidants and pro-antioxidants network: an overview. *Curr Pharm Des* 10: 1677-1694.
- Powell SR (2000) The antioxidant properties of zinc. *J Nutr* 130: S1447-S1454.
- Michalak A (2006) Phenolic compounds and their antioxidant activity in plants growing under heavy metal stress. *Polish J Environ Stud* 15: 523-530.
- Gharras HE (2009) Polyphenols: Food sources, properties and applications-a review. *Int J Food Sci Tech* 44: 2512-2518.
- Bender DA (2003) Vitamin C (ascorbic acid). In: *Nutritional Biochemistry of the Vitamins*. (2nd edn), Cambridge University Press, UK.
- Dawczynski C, Schubert R, Jahreis G (2007) Amino acids, fatty acids, and dietary fibre in edible seaweed products. *Food Chem* 103: 891-899.
- Cornish ML, Garbary DJ (2010) Antioxidants from macroalgae: potential applications in human health and nutrition. *Algae* 25.
- Mabeau S, Fleurence J (1993) Seaweed in food products: Biochemical and nutritional aspects. *Trends Food Sci Technol* 4: 103-107.
- Matanjun P, Mohamed S, Mustapha NM, Muhammad K (2009) Nutrient content of tropical edible seaweeds, *Euclima cottonii*, *Caulerpa lentillifera* and *Sargassum polycystum*. *J Appl Phycol* 21: 1-6.
- Zaytsev O, Cervantes-Duarte R, Montante O, Gallegos-García A (2003) Coastal upwelling activity on the Pacific shelf of the Baja California Peninsula. *J Oceanogr* 59: 489-502.
- Knott KK, Schenk P, Beyerlein S, Boyd D, Ylitalo GM, et al. (2011) Blood-based biomarkers of selenium and thyroid status indicate possible adverse biological effects of mercury and polychlorinated biphenyls in Southern Beaufort Sea polar bears. *Environ Res* 111: 1124-1136.
- Barrera-García A, O'Hara T, Galván-Magaña F, Méndez-Rodríguez LC, Castellini JM, et al. (2012) Oxidative stress indicators and trace elements in the blue shark (*Prionace glauca*) off the east coast of the Mexican Pacific Ocean. *Comp Biochem Physiol C Toxicol pharmacol* 156: 59-66.
- Carvajal M, Martínez MR, Martínez-Sánchez F, Alcaraz CF (1997) Effect of ascorbic acid addition to peppers on paprika quality. *J Sci Food Agric* 75: 442-446.
- Ledezma-Gairaud M (2004) Validación del método: determinación de Vitamina C total por cromatografía líquida de alta resolución. *Tecnología en Marcha* 17: 15-23.
- Singleton VL, Rossi JA (1965) Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J Enol Vitic* 16: 144-158.
- Soriano-Melgar LAA, Alcaraz-Meléndez L, Méndez-Rodríguez LC, Puente ME, Rivera-Cabrear F, et al. (2012) Antioxidant and trace element content of damiana (*Turnera diffusa* Willd) under wild and cultivated conditions in semi-arid zones. *Ind Crops Prod* 37: 321-327.
- Zar JH (1999) *Biostatistical Analysis*. (4th edn), Prentice Hall, Upper Saddle River, NJ, USA.
- Girodon F, Blache D, Monget AL, Lombart M, Brunet-Lecompte P, et al. (1997) Effect of a two-year supplementation with low doses of antioxidant vitamins and/or minerals in elderly subjects on levels of nutrients and antioxidant defense parameters. *J Am Coll Nutr* 16: 357-365.
- Opara EC, Rockway SW (2006) Antioxidants and micronutrients. *Dis Mon* 52: 151-163.
- Gibney MJ, Margetts BM, Kearney JM, Arab L (2004) *Public Health Nutrition*. Wiley, USA.
- Jarup L (2003) Hazards of heavy metal contamination. *Br Med Bull* 68: 167-182.
- Almela C, Algora S, Benito V, Clemente MJ, Devesa V, et al. (2002) Heavy metal, total arsenic, and inorganic arsenic contents of algae food products. *J Agric Food Chem* 50: 918-923.
- van Netten C, Hopton Cann SA, Morley DR, van Netten JP (2000) Elemental and radioactive analysis of commercially available seaweed. *Sci Total Environ* 255: 169-175.
- Kim SY, Sidharthan M, Yoo YH, Lim CY, Joo JH, et al. (2003) Accumulation of heavy metals in Korean marine seaweeds. *Algae* 18: 349-354.
- Besada V, Andrade JM, Schultze F, González JJ (2009) Heavy metals in edible seaweeds commercialised for human consumption. *J Mar Syst* 75: 305-313.
- Almela C, Clemente MJ, Vélez D, Montoro R (2006) Total arsenic, inorganic arsenic, lead and cadmium contents in edible seaweed sold in Spain. *Food Chem Toxicol* 44: 1901-1908.
- Mok JS, Park HY, Kim JH (2005) Trace metal contents and safety evaluation of major edible seaweeds from Korean Coast. *Journal of The Korean Society of Food Science and Nutrition* 34: 1464-1470.
- Mohamed AE, Rashed MN, Moftly A (2003) Assessment of essential and toxic elements in some kinds of vegetables. *Ecotoxicol Environ Saf* 55: 251-260.
- Lawal AO, Audu AA (2011) Analysis of heavy metals found in vegetables from some cultivated irrigated gardens in the Kano metropolis, Nigeria. *J Environ Chem Ecotoxicol* 3: 142-148.
- Muñoz-Barbosa A, Gutiérrez-Galindo EA, Segovia-Zavala JA, Delgadillo-Hinojosa F, Sandoval-Salazar G (2004) Trace metal enrichment in surface sediments of the northwest coast of Baja California, Mexico. *Mar Pollut Bull* 48: 596-603.
- Rodríguez-Meza D, Choumiline E, Méndez L, Acosta-Vargas B, Sapozhnikov D (2007) Chemical composition of sediments and macroalgae Magdalena lagoon complex of Venus. In: Funes R, Gómez J, Palomares R (Eds) *Ecological studies in Magdalena Bay National Polytechnic Institute*.
- McDermid KJ, Stuercke B, Balazs GH (2007) Nutritional composition of marine plants in the diet of the green sea turtle (*Chelonia mydas*) in the Hawaiian Islands. *Bull Mar Sci* 81: 55-71.
- Anantharaman P, Devi GK, Manivannan K, Balasubramanian T (2011)

- Vitamin-C content of some marine macroalgae from Gulf of Mannar Marine Biosphere Reserve, south east coast of India. *Plant Archives* 11: 343-346.
43. Hernández-Carmona G, Carrillo-Domínguez S, Arvizu-Higuera DL, Rodríguez-Montesinos YE, Murillo-Álvarez JI, et al. (2009) Monthly variation in the chemical composition of *Eisenia arborea* J.E. Areschoug. *J Appl Phycol* 21: 607-616.
44. Celikler S, Vatan O, Yildiz G, Bilaloglu R (2009) Evaluation of antioxidative, genotoxic and antigenotoxic potency of *Codium tomentosum* Stackhouse ethanolic extract in human lymphocytes in vitro. *Food Chem Toxicol* 47: 796-801.
45. Vanderslice JT, Higgs DJ, Hayes JM, Block G (1990) Ascorbic acid and dehydroascorbic acid content of foods-as-eaten. *J Food Compos Anal* 3: 105-118.
46. Lim YY, Lim TT, Tee JJ (2007) Antioxidant properties of several tropical fruits: A comparative study. *Food Chem* 103: 1003-1008.
47. Susanna Cunningham- Rundles (1993) Nutrient modulation of the immune response. CRC PressINC.
48. Wintergerst ES, Maggini S, Hornig DH (2006) Immune-enhancing role of vitamin C and zinc and effect on clinical conditions. *Ann Nutr Metab* 50: 85-94.
49. Jiménez-Escrig A, Jiménez-Jiménez I, Pulido R, Saura-Calixto F (2001) Antioxidant activity of fresh and processed edible seaweeds. *J Sci Food Agric* 81: 530-534.
50. Kamiya M, Nishio T, Yokoyama A, Yatsuya K, Nishigaki T, et al. (2010) Seasonal variation of phlorotannin in sargassacean species from the coast of the Sea of Japan. *Phycol Res* 58: 53-61.
51. Erdman JW, Balentine D, Arab L, Beecher G, Dwyer JT, et al. (2007) Flavonoids and heart health: proceedings of the ILSI North America Flavonoids Workshop, May 31-June 1, 2005, Washington, DC. *J Nutr* 137: S718-S737.
52. Williamson G, Manach C (2005) Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr* 81: S243-S255.
53. Brat P, Georgé S, Bellamy A, Du Chaffaut L, Scalbert A, et al. (2006) Daily polyphenol intake in France from fruit and vegetables. *J Nutr* 136: 2368-2373.
54. <http://www.who.int/nutrition/publications/micronutrients/9241546123/en/index.html>.
55. Joint FAO/WHO (1999) Expert committee on food additives. Summary and conclusions, 53rd meeting, Rome, 1-10 June.
56. Joint FAO/WHO (1998) Expert consultation on human vitamin and mineral requirements, Bangkok Thailand, 21-30 September.
57. (1991) Dietary Reference Values for Food Energy and Nutrients for the United Kingdom Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. *Rep Health Soc Subj (Lond)* 41: 1-210.

**Citation:** Tenorio-Rodríguez PA, Méndez-Rodríguez LC, Serviere-Zaragoza E, O'Hara T, Zenteno-Savín T (2013) Antioxidant Substances and Trace Element Content in Macroalgae from a Subtropical Lagoon in the West Coast of the Baja California Peninsula. *Vitam Trace Elem* 2:108. doi:[10.4172/2167-0390.1000108](https://doi.org/10.4172/2167-0390.1000108)

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://omicsgroup.info/editorialtracking/vitamins/>

