

# EFFECT OF *Spirulina platensis* MEAL AS FEED ADDITIVE ON GROWTH, SURVIVAL AND DEVELOPMENT IN *Litopenaeus schmitti* SHRIMP LARVAE

Barbarito Jaime-Ceballos <sup>1\*</sup>, Humberto Villarreal <sup>2</sup>, Tsai Garcia <sup>3</sup>, Lourdes Pérez -Jar <sup>1</sup> and Elvira Alfonso <sup>3</sup>.

(1) Centro de Investigaciones Pesqueras. Ministerio de la Industria Pesquera, 5ta Ave. y 246, Playa, Ciudad Habana. Cuba.

(2) Centro de Investigaciones Biológicas de Noroeste (CIBNOR), Baja California Sur, México.

(3) Centro de Investigaciones Marinas, Universidad de La Habana, Calle 16 No 114, Playa, CP 11300, Ciudad Habana, Cuba.

(\*) Corresponding author: [bjaim04@cibnor.mx](mailto:bjaim04@cibnor.mx), [bjaim04@cip.telemar.cu](mailto:bjaim04@cip.telemar.cu)

## ABSTRACT

To evaluate the effect of *Spirulina platensis* inclusion in microdiets for white shrimp *Litopenaeus schmitti* larvae, a completely randomized experimental design was developed with 4 treatments and three replicates. Diets with 3 inclusion levels (0, 2.5 and 5%) were prepared. A control was fed *Artemia* nauplii. Larvae were fed for 120 hours from mysis I to postlarvae 1 every 4 h, and water quality parameters were measured daily. At the end of the experiment survival was determined, and the development index of the larvae was calculated. The chemical score of the diet was also determined, using the *L. schmitti* postlarvae amino-acid pattern as reference. Tryptophan and histidine were the first and second limiting amino acids in the diets. Survival was around 80% for all treatments. When compared to the control, final postlarval size was significantly smaller. However, the development index (DI) showed that larvae fed diets containing 5% *Spirulina platensis* meal was superior to the rest of the diets and similar to the *Artemia* nauplii ( $p < 0.05$ ).

Key words: shrimp culture; larvae; feed additive; *Litopenaeus schmitti*, *Spirulina platensis*

## RESUMEN

Con el objetivo de evaluar el efecto de la harina de *Spirulina platensis* en dietas microparticuladas para larvas de camarón blanco *Litopenaeus schmitti*, se desarrolló un diseño totalmente al azar con 4 tratamientos y tres réplicas. Se evaluaron tres dietas artificiales (0, 2,5 y 5% de inclusión de harina de *Spirulina*) y un control alimentado con *Artemia* a razón de 5 nauplios.ml<sup>-1</sup>, desde mysis I (M<sub>I</sub>) hasta la primera postlarva (PL<sub>1</sub>), durante 120 horas. Al final del experimento se determinó la talla final y la supervivencia, y se calculó el índice de desarrollo. Se realizó el cómputo químico de las dietas, tomando como referencia el patrón aminoacídico de las postlarvas de *L. schmitti*. Triptofano e histidina fueron el primer y segundo aminoácidos limitantes en todos los tratamientos. La supervivencia se mantuvo cerca del 80% para todos los casos. El crecimiento de las larvas no presentó diferencias significativas ( $p > 0.05$ ) entre los alimentos microparticulados ensayados, obteniéndose la talla final mayor al utilizar nauplios de *Artemia*. Las larvas que consumieron el alimento vivo y la dieta microparticulada enriquecida con 5% de *Spirulina* alcanzaron los mayores valores de índice de desarrollo.

Palabras clave: cultivo decamarones; larvas; aditivo alimenticio; *Litopenaeus schmitti*, *Spirulina platensis*.

Changes in abundance and distribution of wild penaeid postlarvae and the increasing world market demand for prawns has led to rapid development of commercial larval breeding systems for penaeids. (Hudinaga and Miyamura, 1962).

Penaeid shrimp hatcheries are highly dependent on live phytoplankton and zooplankton cultures, which are costly and labor intensive (Robinson *et al.*, 2005). However, recent advances in formulation and microparticulate diet technology will soon permit the completion of the larval cycle exclusively on inert diets (Le Moullac *et al.*, 1987; Kanazawa 1990; Wouters *et al.*, 2004).

The search of efficient feeds for the white shrimp larvae is necessary in order to increase postlarvae quality and culture profitability. The dependence on *Artemia* nauplii for production is a concern for commercial hatcheries due to the possibility of a global shortage and the rising cost of *Artemia* cyst, as well as the possible variations in nutritional value and the risk of disease transfer. Over 80% of hatcheries reported decreasing *Artemia* cyst utilization in the last few years as a result of these concerns (Wouters *et al.*, 2003).

One solution to this problem is to develop a practical and economical formulation capable of satisfying the nutritional requirement of the different larval stages.

In Cuba, commercial postlarvae production of *Litopenaeus schmitti* still depends on live food such as *Artemia* cysts and microalgae (*Chaetoceros* spp. and *Tetraselmis* spp.), as well as on imported microparticulated feeds in order to guarantee the post larvae demands from the shrimp farms.

As feed additives, dried algae improve growth, feed efficiency, carcass quality, and physiological response to stress and disease in several species of fish (Mustafa and Nakagawa, 1995). Experimental formulations for white shrimp larvae have also been successful (Jaime *et al.*, 1996; Márquez 1997; Artiles *et al.*, 1999).

*Spirulina* is considered a rich source of protein, vitamins, minerals, essential amino acids, and fatty acids (gamma - linolenic acid (GLA), and antioxidant pigments, such as carotenoids (Belay *et al.*, 1996). In addition, it is effective as an immunomodulator (Takeuchi *et al.*, 2002). Several studies have been conducted using dried *Spirulina* as a feed supplement (Chow and Woo, 1990; Watanabe *et al.*, 1990), and *Spirulina platensis*, which is commercially produced in Cuba, has been considered for partial substitution of microalgae in the feeding of white shrimp *Litopenaeus schmitti* protozoas (Jaime, *et al.*, 2004).

The present study evaluates the effect of *Spirulina platensis* meal, as feed additive, on body length, survival and development index in *L. schmitti* larvae.

## MATERIALS AND METHODS

Three artificial diets (0, 2.5 and 5% of *Spirulina* meal) and a control (*Artemia* fed at 5 nauplii. ml<sup>-1</sup>) were tested in a totally randomized design during 120 hours, using *Litopenaeus schmitti* (Pérez-Farfante and Kensley, 1997) larvae obtained from a commercial rearing tank of the laboratory of postlarvae production "YAGUACAM", in Cienfuegos, Cuba. Mysis I larvae were stocked at a density of 120 organisms per liter, in 12 fiber-glass tanks, using three replicates per diet, with 100 liters of filtered (5 µm) and uv sterilized seawater at 35 ups. Constant aeration was provided via air stones using a 5 hp turbo blower. Na<sub>2</sub> EDTA was added to the water at 10 mg.l<sup>-1</sup> (Lawrence *et al.*, 1981). To eliminate faecal residues and food remains, a 50% water exchange was done daily. Constant illumination was provided with fluorescent lights, temperature (°C), oxygen (mg.l<sup>-1</sup>), salinity (psu), pH and ammonium (µg.l<sup>-1</sup>) were registered daily before the water exchange.

Three microparticulated diets were elaborated (Table 1), using the MIP-CIP 2 formulation evaluated by Artiles *et al.*, (1999) as reference. Particle size ranged from 100 to 250 µm. Feeding was carried out every 4 hours (06.00, 10.00, 14.00, 18.00, 22. 00 and 02.00 hours) at a rate of 0.3 mg/larvae/day of food. Proximal composition of diets was determined according to the procedures of AOAC (1995). Total pigments were determined using the trichromatic method (Strickland and Parson, 1972).

The chemical score of the experimental diets was carried out using the amino acid composition of *L. schmitti* postlarvae (Gallardo *et al.*, 1989) as reference.

Survival was determined using the volumetric method, taking five 100 ml samples per tank daily. The number of mysis and postlarvae in each sample was determined in order to calculate the development index (DI) following Villegas and Kanazawa (1979). Larval growth was determined at the end of the study. Using an optical microscope (8x), 40 animals were measured per tank from the tip of the rostrum the end of the tail, excluding the thorns.

ANOVA and Duncan's multiple range tests were used to analyze the final data after using Kolmogorov-Smirnov and Bartlett tests to confirm homoscedasticity (Sigarroa, 1985).

## RESULTS AND DISCUSSION

Water quality parameters were maintained within the range recommended by Vega and de la Cruz (1988) for the species. Dissolved oxygen was  $7.25 \pm 0.25$  mg.l<sup>-1</sup>, temperature was maintained at  $27.6 \pm 1^\circ\text{C}$ , and pH was  $8.05 \pm 0.25$ . NH<sub>4</sub>-N had a range between 328.1- 736.3 µg.l<sup>-1</sup>.

Mean body length, survival and the DI are presented in Table 2. Larvae fed the *Artemia* nauplii control were significantly bigger ( $p < 0.05$ ). There were no significant differences between treatments for larval size when fed artificial diets. The results are similar those of Artiles *et al.*, (1999).

Jones (1998) indicated that only when a formulated artificial diet is accepted, ingested and assimilated at rates comparable to live feeds will it be possible to investigate specific nutritional requirements. Consequently, the size, shape, attractiveness and nutritional content of an

Table 1. Proximal composition of experimental diets.

Ingredients (%)	Dietary group		
	SP0	SP2.5	SP5.0
Fish meal <sup>1</sup>	35	32.5	30
Soybean meal <sup>3</sup>	13	13	13
Shrimp meal <sup>4</sup>	15	15	15
Squid meal <sup>4</sup>	16	16	16
Spirulina meal <sup>2</sup>	-	2.5	5
Fish oil	4.5	4.5	4.5
Vegetable oil	4.5	4.5	4.5
Premix Vit + Min <sup>5</sup>	5	5	5
Binder CMC*	5	5	5
Vit.C	0.5	0.5	0.5
Cholesterol	0.5	0.5	0.5
Total	100	100	100
<b>PROXIMATE COMPOSITION (%)</b>			
Moisture	8.4 ± 0.06	8.1 ± 0.05	7.9 ± 0.07
Crude protein	46 ± 0.08	46.9 ± 0.12	47 ± 0.09
Lipid	13.9 ± 0.04	14.2 ± 0.06	14.4 ± 0.02
Ash	8.3 ± 0.02	8.4 ± 0.04	8.5 ± 0.03
Total pigments	0.05 ± 0.003	0.48 ± 0.07	0.84 ± 0.08
Crude Fibber	2.5 ± 0.02	2.6 ± 0.04	2.7 ± 0.05

\* Carboxymethylcellulose sodium

<sup>1</sup> Anchoveta fish meal (*Engraulis* sp.) CORPESCA S.A. Chile.

<sup>2</sup> *Spirulina platensis*. GENIX, La Habana, Cuba.

<sup>3</sup> Molinera de Santiago de Cuba. Cuba

<sup>4</sup> Shrimp meal (*Litopenaeus schmitti* y *Farfantepenaeus notialis*) and Squid meal (*Loligo* sp.), made in laboratory.

<sup>5</sup> Formula of the Union of Companies feeds MINAGRI. Cuba, for commercial feeds for shrimp culture. One metric ton of premix.

Retinol 12.500. UI; Thiamine 10.000 mg; Riboflavin 20.000, mg; Pyridoxine 10.000 mg; Cianocobalamin 40,0 mg; Ascorbic acid 500.000 mg; Cholecalciferol 0,5 2.400.000 UI; DL- $\alpha$ -tocopherol 100.000; Menadione 4.000 mg; Pantothenic acid 40.000 mg; Choline chloride 1.600. mg; Folic acid 2.000 mg; Nicotinic acid , 140.000 mg; Biotin 1.000 mg; Inositol 300.000 mg; Paraminobenzoic acid 35.000 mg; Cobalt 200 mg; Copper 2.000 mg; Iron 20.000 mg; Iodine 1.500 mg; Manganese 40.000 mg; Zinc 20.000 mg; Selenium 100,0 mg

Table 2. Mean  $\pm$  S.D. for final length, survival and development index (DI) of *L. schmitti* larvae fed diets with different *Spirulina platensis* inclusion levels\*.

Treatment	Final Length (mm)	Survival (%)	Development Index
SP0	3.43 $\pm$ 0.02 <sup>a</sup>	86.1 $\pm$ 4.3 <sup>a</sup>	6.72 $\pm$ 0.03 <sup>a</sup>
SP2.5	3.48 $\pm$ 0.05 <sup>a</sup>	80.5 $\pm$ 3.8 <sup>a</sup>	6.78 $\pm$ 0.04 <sup>b</sup>
SP5.0	3.44 $\pm$ 0.03 <sup>a</sup>	84.2 $\pm$ 3.5 <sup>a</sup>	6.82 $\pm$ 0.04 <sup>c</sup>
Control	3.71 $\pm$ 0.03 <sup>b</sup>	85.3 $\pm$ 3.1 <sup>a</sup>	7.00 $\pm$ 0.12 <sup>c</sup>

\* Columns with the same letter are not significantly different ( $p > 0.05$ ).

artificial diet must be optimized so that rates of ingestion result in growths and survivals similar to those achieved with live feeds. In the present study, the results when feeding artificial diets resulted in similar rates of survival compared to the control. Kumlu and Jones (1995) reported similar survivals for *Farfantepenaeus indicus* when feeding microparticulated diets. In this case, mortality was probably associated to handling during the water exchange process. This is in agreement with Alfonso (1996), who stated that, when the quality of food is good, the survival is determined by manipulation.

Arellano *et al.*, (1993) feeding *Litopenaeus vannamei* mysis with microparticulated diets, achieved smaller growth rates and survival that when they combined the artificial diet with *Artemia* nauplii, suggesting that these constitute an important source of nourishment in this phase. However, Kurmaly *et al.*, (1989) achieved the metamorphosis from M<sub>II</sub> - PL1 of *Penaeus monodon*, feeding a microencapsulated diet rich in high quality proteins and lipids demonstrating the possibility to completely replace live food during the larval development with artificial rations, whenever these meet the nutritional requirements for the species.

Márquez (1997) testing diets with different levels of torula yeast (*Sacharomyces cereviseae*) with *L. schmitti* achieved similar final postlarvae sizes to the ones obtained in this work. The growth results indicate that supplementation of *S. platensis* meal did not favor an increment in size of the shrimp.

Jaime *et al.*, (1996) evaluating different commercial microparticles for *L. schmitti* larvae found that better growth rates were obtained with diets containing 17% lipids, when compared with experimental diets containing 8.6% lipids. Artiles *et al.*, (1999), on the other hand, obtained the best growth results when combining *Artemia* nauplii with a microdiet containing 14% lipids. This formula was used in this work as control. The results show that, for the formulated diets, the inclusion of *S. platensis* did not affect survival or final size.

Kumlu and Jones (1995) reported delays of up to 2 days in larval development, of *Fenneropenaeus indicus* when fed with microencapsulated diets. On the other hand, Artiles *et al.*, (1999) and Márquez (1997) report that *L. schmitti* reach the PL<sub>1</sub> stage after 120 hours when fed with a combination of microparticulated diet and *Artemia* nauplii. The time of development was similar for this trial. However, the rate of metamorphosis, expressed through the development index (DI) was statistically different for the four treatments, indicating that the concentration of *Spirulina platensis* influenced development, and that an inclusion rate of 5%, DI was not statistically different to that of the *Artemia* control. Further studies should incorporate higher inclusion levels of *S. platensis* in the diet so as to assess the potential to increase final size, and the potential for partial substitutions of the *Artemia* nauplii.

The determination of nutrient essentiality to larval penaeid shrimp is complicated by a life cycle consisting of a series of planktotrophic stages resulting in substantial trophic level changes (Jones *et al.*, 1993). Cuzon (2004) states that there is a relationship between whole body composition, in terms of essential amino acids for *Penaeus monodon* juveniles, and its protein requirement. So, we suggest that the chemical score can be considered an index to evaluate the protein quality of feeds for aquatic species. Fig. 1 shows that the formulations of the experimental diets are similar in terms of amino acid balance. However, the chemical score shows that the experimental diets are limiting for several amino acids, when compared with *L. schmitti* postlarvae. Hysitidine and Thriptophan are the first limiting amino acids.

This is similar to the report of Gaxiola *et al.*, (2002) for *Litopenaeus setiferus* and *Litopenaeus vannamei*. The reduction in final size could be related to this, and further research should evaluate microparticulated diets that supply all of the essential amino acids.

Table 1 shows the level of total pigments in the diets. It appears that as the level of *S. platensis* increased in the diet and the pigment concentration increases, the physiological and metabolic responses of *L. schmitti* larvae improves. Several authors mention that this microalgae has an important role in metabolic, antioxidant, respiratory and immunological roles of aquatic organisms (Mustafa *et al.*, 1994; Miyasaki *et al.*, 1995; Gabaudan 1998; Takeuchi *et al.*, 2002).

This is in agreement with the report by Meyer (2000) who suggests that diets for aquatic larvae must be formulated to provide suitable amounts of supplemental carotenoids, specifically astaxanthin, which cannot be synthesized “*the novo*”, but depends on the assimilation of a precursor Ringelberg (1980).

It is important to indicate that the use of microparticles for shrimp larvae production would represent a significant decline in production costs, particularly due to the reduction or elimination of areas required for supporting cultures, such as microalgae and *Artemia*. This use could also provide advantages in terms of disease and water quality control, and the possibility of recycling the water, further improving the operational and practical efficiency of the culture (Robinson 2005).

## CONCLUSION

- The development index showed that *Litopenaeus schmitti* larvae fed diets containing 5% *S. platensis* meal produced results similar to those obtained by feeding *Artemia* nauplii.
- Final length were significantly smaller when feeding microparticles, which may be related to nutritional deficiencies, particularly associated to essential amino acids.
- Further improvement in formulation and management of rearing tanks should lead to the full replacement of live food by inert particles in commercial hatcheries for the species.

## ACKNOWLEDGEMENTS

The authors are grateful to the staff at Centro de Investigaciones Pesqueras, YAGUACAM hatchery and Centro de Investigaciones Marinas (Cuba).

Barbarito Jaime is a doctoral fellow of CONACYT (No. 182852) at the Centro de Investigaciones Biológicas del Noroeste (México).




Fig. 1. Chemical score (%) of the experimental diets in relation to the amino acid composition of *L. schmitti*.

**REFERENCES**

Alfonso, E. (1996): Larvicultura de Camaroes Marinhos. In: *Anais do Primeiro Workshop do Estado do Ceará sobre Cultivo de Camarao mariho*. (T.C.V. Gesteira and A.J.P. Nunes, eds). Fortaleza, Ceará, Brasil. pp. 86-100

AOAC, (1995): *Official Methods of Analysis of the Association of Official Analytical Chemist*. Vol. I, 16th ed. Washington, D. C., USA. 1234 p.

Arellano, E.M., F. Carvaca, L. Gómez y A. Pedrazzoli (1993): Proyecto optimización del cultivo de larvas de camarón, "ocular". In: *Memorias de de Edgar Arellano. Once años dedicados a la investigación y desarrollo de la Acuicultura*. ESPOL-CENAIM. 1993 San Pedro de Manglaralto, Ecuador. pp, 19-24.

Artiles, A.M., I. Fraga, J. Galindo y B. Jaime (1999): Empleo de dietas artificiales en la larvicultura del camarón blanco *Penaeus schmitti*. *Rev. Invest. Mar.* 20(1-3): 75-81.

Belay, A., T. Kato y Y. Ota (1996): *Spirulina (Arthrospira)*: Potential application as an animal feed supplement. *J. Appl. Phycol.* 8: 303-311.

Chow, C.Y. y N.Y.S. Woo (1990): Bioenergetics studies on an omnivorous fish *Oreochromis mossambicus*: Evaluation of the utilization of Spirulina algae in feed. In: *Proceeding of the 2nd Asian Fisheries Forum* (R.Hirano and I. Hanyu, eds). The Asian Fisheries Society; Manila, pp: 291-294.

Cuzon. G., J. Guillaume y G. Gaxiola (2004): Review on amino acid requirement in shrimp. (Abstract Book) *Aquaculture. 2004*. Marzo 1-5, 2004. Hawaii Convention Center. Honolulu, Hawaii, USA. 140 pp.

Gabaudan, J. (1998): Dietary astaxanthin improves production yield in shrimp farming. *Fish-Chimes*, 18: 37-39.

Gallardo, N., R. González, O. Carrillo, O. Valdés y A. Forrellat (1989): Una aproximación a los requerimientos de aminoácidos esenciales de *Penaeus schmitti*. *Rev. Invest. Mar.* 10(3):259-267.

- Gaxiola G., P. Gallardo, R. Ravallec, C. Durruty, T. García, G. Cuzon, A. Van Wormhoudt y A. Pedroza (2002): Avances en el uso de alimentos artificiales en la larvicultura de camarón.. In: *Avances en Nutrición Acuícola VI. Memorias del VI Simposium Internacional de Nutrición Acuícola*. (L. E. Cruz-Suárez, D. Ricque-Marie, M. Tapia-Salazar, M.G Gaxiola-Cortés and N. Simoes, eds.). 3 al 6 de Septiembre del 2002. Cancún, Quintana Roo, México. pp: 227-242
- Hudinaga, M. y M. Miyamura (1962): Breeding of the kuruma prawn (*Penaeus japonicus* Bate). *Journal Oceanogr. Soc. Jpn.* (20<sup>th</sup> Anniv. Vol ): 694-706.
- Jaime, B., A.M. Artilles, J. Galindo e I. Fraga (1996): Evaluación de alimentos microparticulados en el cultivo larval de *Penaeus schmitti*. *Rev. Cub. Invest. Pesq.* 20(2):22-24.
- Jaime-Ceballos, B., H. Villarreal-Colmenares, T. García-Galano, R. Civera-Cerecedo y G. Gaxiola-Cortés (2004): Empleo del polvo de *Spirulina platensis* en la alimentación de zoeas y mysis de *Litopenaeus schmitti* (Pérez-Farfante and Kensley, 1997). In: *Avances en Nutrición Acuícola VII, Memorias del VII Simposium Internacional de Nutrición Acuícola*. (L. E. Cruz-Suárez, D. Ricque Marie, M. G. Nieto López, D. Villarreal, U. Scholz and M. González, eds) 16-19 Nov, 2004, Hermosillo, Sonora, México, pp: 617-635.
- Jones, D.A. (1998): Crustacean Larval microparticulaed diets, *Review in Fisheries Science*, 6(1-2): 41-54.
- Jones, D.A., M.S. Kamarudin y L. LeVay (1993): The potential replacement of live feed in larval culture. *Journal of the World Aquaculture Society* 24(2):199-210.
- Kanazawa, A. (1990): Microparticulate feeds for Penaeid larvae. In: *Advances in tropical aquaculture* (Barred, J. ed.) Workshop held in Tahiti (French Polynesia), February 20-March 4, 1989. 1990 (9): 395-404.
- Kumlu, M. y D.A. Jones (1995): The effect of live and artificial diets on growth, survival and trypsin activity in larvae *Penaeus indicus*. *Journal of the World Aquaculture Society* 26(4):406-415.
- Kurmaly, K., D.A. Jones, A.B. Vule y J. East (1989): Comparative analysis of the growth and survival of *Penaeus monodon* larvae, from protozoa 1 to postlarvae 1 a live feeds, artificial diets and a combination of both. *Aquaculture*, 81: 27-45.
- Lawrence, A., J. Fox y F. Castille (1981): Decreased toxicity of cooper and manganese ions to shrimp nauplii (*Penaeus stylirostris*, Simpson) in the presence of EDTA. *World Mariculture Society*. 41: 1-18.
- Le Moullac, L., R. Leguedes, G. Cuzon y J. Goguenheim (1987): *Microparticulate diets for penaeid shrimp larvae*. AQUACOP, IFREMER. WAS. Guayaquil, Ecuador, 1987
- Márquez, C.G. (1997): *Evaluación de la levadura torula en la alimentación de larvas y postlarvas de camarón blanco Penaeus schmitti, con dietas artificiales*. Tesis de Maestría, Centro de Investigaciones Marinas, Universidad de La Habana, 60 pp.
- Meyers, S.P. (2000): Papel del carotenoide astaxantina en nutrición de especies acuáticas. In: *Avances en Nutrición Acuícola IV. Memorias del IV Simposium Internacional de Nutrición Acuícola*. (R. Civera-Cerecedo, C. J. Pérez-Estrada, D. Ricque-Marie, and L. E. Cruz-Suárez, eds.) Noviembre 15-18, 1998. La Paz, B.C.S., México. pp: 473-491
- Miyasaki, T., M. Sato, R. Yoshinaka y M. Sakaguchi (1995): Effect of vitamin C on lipid and carnitine metabolism in rainbow trout . *Fish. Sci* 61, 501-506.
- Mustafa, M.G. y H. Nakagawa (1995): A review: Dietary benefits of algae as an additive in fish feed. *Isr. J. Aquacult.* 47, 155-162.
- Mustafa, M.G., T. Umino y H. Nakagawa (1994): The effect of *Spirulina* feeding on muscle protein deposition in red sea bream, *Pagrus major*. *J. Appl. Ichthyol.* 10, 141-145.
- Pérez-Farfante, I. y B. Kensley (1997): Penaeoid and sergestoid shrimps and prawns of the world, keys and diagnoses for the families and genera. *Mémoires Muséum National d' Histoire Naturelle, Zoologie*, 175, 235.
- Ringelberg, J. (1980): Aspects of red pigmentation in zooplankton, especially copepods. In: *Evolution Ecology of Zooplankton Communities* (W.C Kerfoot., ed.). Univ. of New Hampshire, Hanover. pp. 91-97.
- Robinson, C.B., T.M. Samocha, J.M. Fox, G. Gandy y D.A. Mckee (2005): The use of inert artificial commercial food sources as replacements

of traditional live food items in the culture of larval shrimp. *Farfantepenaeus aztecus*. *Aquaculture*. 245:135-147.

Sigarroa, A. (1985): *Manual de prácticas de biometría y diseño experimental*. Editorial Pueblo y Educación. Ciudad de La Habana. 793 pp.

Strickland J.D.H. y T.R. Parsons (1972): *A practical handbook of seawater analysis*. *Fisheries Research Board of Canada Bulletin*, 310 pp.

Takeuchi, T., J. Lu, G. Yoshizaki y S. Satoh (2002): Effect on the growth and body composition of juvenile tilapia *Oreochromis niloticus* fed raw *Spirulina*. *Fisheries Science*. 68:34-40

Vega, A.J. and A. de la Cruz (1988): Influencia de la temperatura, la salinidad y el pH sobre las larvas del camarón blanco *Penaeus schmitti*. *Rev. Invest. Mar.* 9(1):25-38.

Villegas, D.K. y A. Kanazawa (1979): Relationship between diet and growth of zoeal and mysis stages of *Penaeus japonicus* Bate. *J. Zool.* 10(2):305-399.

Watanabe, T., W. Liao, T. Takeuchi y H. Yamamoto (1990): Effect of dietary *Spirulina* supplement on growth performance and flesh lipids of cultured striped jack. *J. Tokyo Univ. Fish.* 77: 231-239.

Wouters, R., T. Van Horenbeek, G. Merchie y P. Bribson (2004): Dietas larvales para camarón. Dietas secas elaboradas con ingredientes marinos frescos. *Panorama Acuicola Magazine* Mar/Abr, 9, 54-55.

Wouters, R., T. VanHorenbeek y G. Merchie (2003): Flakes for shrimp postlarvae nutrient, retention and shrimp culture performance. *Panorama Acuicola Magazine*. Mar/Abr. 8(3):52-53.

Aceptado: 15 de octubre del 2005